

OFFPRINT

RGZM – TAGUNGEN Band 40

Sonja B. Grimm · Mara-Julia Weber
Ludovic Mevel · Iwona Sobkowiak-Tabaka (eds)

FROM THE ATLANTIC TO BEYOND THE BUG RIVER

FINDING AND DEFINING THE FEDERMESSER-GRUPPEN / AZILIAN

Proceedings of Session A5b (Commission »The Final Palaeolithic of
Northern Eurasia«) of the XVIIth UISPP Congress,
Burgos, September 2014

Gedruckt mit Unterstützung des
Zentrums für Baltische und Skandinavische Archäologie
in der Stiftung Schleswig-Holsteinische Landesmuseen
Schloss Gottorf, Schleswig



**Römisch-Germanisches
Zentralmuseum**
Leibniz-Forschungsinstitut
für Archäologie



Redaktion: Gundula Lidke (ZBSA) unter Mitwirkung von Sonja B. Grimm
und Mara-Julia Weber (beide ZBSA); Claudia Nickel (RGZM)
Satz: Claudia Nickel (RGZM)
Umschlaggestaltung: Claudia Nickel (RGZM)

Bibliografische Information der Deutschen Nationalbibliothek

Die Deutsche Nationalbibliothek verzeichnet diese Publikation in
der Deutschen Nationalbibliografie; detaillierte bibliografische
Daten sind im Internet über <http://dnb.d-nb.de> abrufbar.



Dieses Werk ist unter der Creative Commons-
Lizenz 4.0 (CC BY-NC-ND 4.0) veröffentlicht.

Propylaeum
FACHINFORMATIONSDIENST
ALTERTUMSWISSENSCHAFTEN

Publiziert bei Propylaeum, Universitätsbibliothek Heidelberg /
Römisch-Germanisches Zentralmuseum, Mainz 2020.

© 2020 Verlag des Römisch-Germanischen Zentralmuseums, Mainz

Diese Publikation ist auf <http://www.propylaeum.de> dauerhaft frei
verfügbar (Open Access).

URN: [urn:nbn:de:bsz:16-propylaeum-ebook-575-4](http://nbn:de:bsz:16-propylaeum-ebook-575-4)

DOI: <https://doi.org/10.11588/propylaeum.575>

e-ISBN: 978-3-948465-04-9 (PDF)

e-ISSN: 2698-9069

ISBN: 978-3-948465-05-6 (Softcover)

ISSN: 1862-4812

CONTENTS

Sonja B. Grimm · Mara-Julia Weber · Ludovic Mevel · Iwona Sobkowiak-Tabaka

Not Quite as Far as from the Atlantic to beyond the Bug River – an Editorial VII

William Mills

Is there any Order to this Chaos? Patterns of Space, Stone, and Place
in Lateglacial South-East England and the Channel River Network 1

Olivier Bignon-Lau

About the Early Azilian Way of Life in the Paris Basin: Economical and Spatial Insights
from Zooarchaeological Data 25

Philippe Crombé

The Environmental Setting for the Lateglacial Recolonisation of the
Scheldt Basin (North-West Belgium) by the Federmesser-Gruppen 51

Florian Sauer

Landform-Based Modelling of Potential Biological Diversity. Inferring Ecological Variability
in Site Catchments from Digital Elevation Data 67

Martin Moník · Anna Pankowská

Settlement Patterns of the Late Palaeolithic in Bohemia and Moravia 79

Katarzyna Pyżewicz · Witold Grużdź · Piotr Rozbiegalski · Aleksandra Rakoca

Two Methods of Blade Production among Arch-Backed Point Groups –
a Case Study from the Western Polish Lowland 91

Paweł Valde-Nowak · Anna Kraszewska

Two Late Palaeolithic Arch-Backed Points Varieties in the Northern Carpathians 105

Iwona Sobkowiak-Tabaka

Similar, yet Different. The Arch-Backed Piece Technocomplex in Poland 119

List of Contributors 139

NOT QUITE AS FAR AS FROM THE ATLANTIC TO BEYOND THE BUG RIVER – AN EDITORIAL

To Martin Street, our friend and teacher

The history of this volume begins in September 2012 when three of the editors met at MONREPOS Archaeological Research Centre and Museum for Human Behavioural Evolution (part of the Römisch-Germanisches Zentralmuseum, Leibniz-Forschungsinstitut für Archäologie, a member of the Leibniz Association). While Sonja B. Grimm has been writing her PhD about the transition from Magdalenian to Federmesser-Gruppen in North-West Europe (Grimm 2019) there, Ludovic Mevel stayed for several months to analyse some of these assemblages technologically to compare them to French inventories within a Post-Doc project (Mevel/Grimm 2019). During this time, Iwona Sobkowiak-Tabaka came to examine some Federmesser-Gruppen material from Central Rhineland sites for comparative reasons within her habilitation project about the Federmesser-Gruppen on the North European Plain (Sobkowiak-Tabaka 2017). She was hosted by Martin Street who also functioned as Sonja's tutor. In this culture clash of English, French, German, and Polish traditions of thoughts on the Federmesser-Gruppen and Azilian, we realised that it could be very fruitful to bring together these different approaches of material culture and spatial studies. In a direct comparison we hoped to find answers to questions such as:

What do we actually know about the groups living in Western and Central Europe during the Weichselian Lateglacial Interstadial? How are they reflected in the archaeologically defined groups that we call Azilian and Federmesser-Gruppen? Do they represent related groups or even a single entity? Or were they formed by many rather different groups of hunter-gatherers with a similar way of life? Were they contemporaries or chronological successors? And how are these related to the Epigravettian industries in Southern and South-eastern Europe?

In order to gather different ways of approaching these questions, Martin Street suggested to organise a session on behalf of the UISPP commission »The Final Palaeolithic of Northern Eurasia« but left the details to us »younger ones«. Yet, Martin's influence as researcher, mentor, host, peer, and friend was always present during the process of organising and holding the session as well as in the production of this volume.

Considering the geographic extent of assemblages attributed to the Azilian and/or Federmesser-Gruppen and the differences therein, we quickly dropped the idea of also including the Epigravettian. Instead, to include directly the Northern European research tradition in the organising team, we turned to Mara-Julia Weber working in Schleswig and together prepared a session for the XVIIth UISPP congress. So in September 2014, a session with the title »From the Atlantic to beyond the Bug River – Finding and defining the Federmesser-Gruppen/Azilian on the North European Plain and adjacent areas« was held in Burgos (Spain). It consisted of an introductory talk, nine presentations and a poster covering the area from the United Kingdom, France, Germany, the Czech Republic, and Poland. Hence, from the northern Carpathians to beyond the English Channel would have been the more appropriate title to describe the actual geography covered by our contributions. Three of these presentations were – by the time – on-going PhD projects (W. Mills, M. Monik, F. Sauer) which showed that the questions relate to a topical subject of study and the results will be picked up by a young generation of scientists.

However, the variety of nomenclature already displayed in these talks showed the necessity but also the difficulty of the prospective discussion. Federmesser-Gruppen and Azilian were used almost evenly but also Arch-Backed Point (ABP) groups, Lateglacial, Late Palaeolithic, and Late Upper Palaeolithic were used to describe the material from this period. This plurality might indicate a few different groups but often they rather referred to different scales or appeared rudiments of different scholarly traditions (cf. Sauer/Riede 2019). In general, terms such as Federmesser-Gruppen, Azilian, Penknife Point phase, Tjongerian, Curve-Backed Point groups, or ABP are general representatives for a period when foragers roamed in a boreal environment and, thus, were already different from the classic Upper Palaeolithic hunter-gatherers of the Late Pleistocene steppe landscapes but were not fully Mesolithic yet. The difference to perhaps more local groups such as the Tjonger group, Witowian, or Atzenhof group is often fuzzy. These variable levels reflect different spatial and chronological sections of the hardly distinguishable scale of human social units. It is generally difficult to identify these units based on archaeological material that is then compared over increasingly large areas and timespans. This taxonomic approach becomes particularly difficult if it is applied to periods when only low cost strategies are used (Vaquero/Romagnoli 2018) such as it appears during the here discussed period. The decrease of elaborate and standardised behaviours – compared to the previous Magdalenian technological behaviours – reflected in the material record give only very few possibilities to link differences in the material to group traditions. These few and/or small differences are then likely to be overestimated.

Besides the difficulty of identifying the material, processes during and after the Younger Dryas also affected the *in situ* preservation and visibility of the material. For instance, taphonomic disturbances of the material deposition occurred when deep frozen grounds melted and caused significant *in situ* sediment loss or the material became inaccessible due to cover by sometimes very thick coversands. Finally, the preservation conditions become less favourable for organic material in environments with richer biogenic interactions and reduced mineral coverage. The long bog development that occurred during the Holocene seemed to have no equivalent in the Lateglacial Interstadial. Hence, less diverse remains were preserved to base the studies on.

Furthermore, chronostratigraphic terms such as Meiendorf, Bølling, or Allerød were also used differently (cf. Terberger/Barton/Street 2009, tab. 1). So when trying to at least correlate the archaeological terms spatially and chronologically again quite some confusion can be caused. This also applies to the radiocarbon dates that are sometimes given as raw radiocarbon data (in this volume identified as ¹⁴C-BP) and sometimes calibrated data (in this volume given as cal. BC). The difference is occasionally not visible.

Hence, before being able to explore the inconsistent use of terminology as one of the main aims of the session, we had to find a way between the wide-spread wish to remain within the own frame of traditions and the establishment of a shared frame of references. Although we attempted to unify some general terminology within the different contributions, different concepts remained and the need for a common analysis to come up with a classification we can all work with is still a desideratum. However, single-minded attempts will not solve the problem but rather add yet another new taxonomy that has not grown from a reunification of traditions. Instead of this top-down approach, a bottom-up approach of unifying the recording systems is on its way, for example the work of L. Mevel using one technological approach to describe the assemblages from the Paris Basin and the Rhineland was taken to the north in cooperation with M.-J. Weber and will be continued in a PhD thesis about the Federmesser-Gruppen in northern Germany and Denmark by T. Burau. Anyway, to come to results that can be compared across wider geographic areas, we all have to leave the comfort zone of our national narratives and analytical traditions and see what in a greater framework is possible. In this volume, we assemble contributions from different parts of North-Western and Central Europe (France, UK, Belgium, Germany, Poland, Czech Republic). Lithic industries still remain the main archaeological source (Sobkowiak-Tabaka; Pyżewicz et al.; Valde-Nowak and Kraszewska) but recent research has also been

exploring and combining further sources of information such as faunal and spatial data and geo- and environmental archives (Bignon-Lau; Mills; Crombé; Sauer; Monik and Pankowská). With regard to the research topics addressed, one can observe two main perspectives by which the relationship between archaeological entities in the mid-Lateglacial Interstadial is regarded: spatial behaviour on variable scales, from site to landscape, on the one hand and the cultural attribution of single lithic inventories on the other hand.

We may not have come closer in defining the Federmesser-Gruppen and/or Azilian but we hope that this book will keep the discussion about these Final Palaeolithic hunter-gatherer groups on-going, especially as they received some relevance in relation to the Epigravettian in the light of aDNA results (Posth et al. 2016; Fu et al. 2016).

Acknowledgements

The authors want to thank the former board of the UISPP commission »The Final Palaeolithic of Northern Eurasia« (Berit Valentin Eriksen, Eelco Rensink, and Martin Street) for their support of the idea of the session and the publication. We also like to thank the RGZM for accepting these proceedings for publication. Furthermore, we wish to express our deep gratitude to the Centre of Baltic and Scandinavian Archaeology (ZBSA) for the generous financial support during the editing and publishing process. This support made the employment of Dr. Gundula Lidke possible to whom we owe a big thank you for her comprehensive text and picture editing and her kind communications with the authors of the different contributions. We also much appreciated the time donated to this project by the peer reviewers and English proof-readers who clearly improved the science presentation in this volume.

References

- Fu et al. 2016: Q. Fu / C. Posth / M. Hajdinjak / M. Petr / S. Mallick / D. Fernandes / A. Furtwängler / W. Haak / M. Meyer / A. Mittnik / B. Nickel / A. Peltzer / N. Rohland / V. Slon / S. Talamo / I. Lazaridis / M. Lipson / I. Mathieson / S. Schiffels / P. Skoglund / A. P. Derevianko / N. Drozdov / V. Slavinsky / A. Tsybankov / R. G. Cremonesi / F. Mallegni / B. Gély / E. Vacca / M. R. G. Morales / L. G. Straus / C. Neugebauer-Maresch / M. Teschler-Nicola / S. Constantin / O. T. Moldovan / S. Benazzi / M. Peresani / D. Coppola / M. Lari / S. Ricci / A. Ronchitelli / F. Valentin / C. Thevenet / K. Wehrberger / D. Grigorescu / H. Rougier / I. Crevecoeur / D. Flas / P. Semal / M. A. Mannino / C. Cupillard / H. Bocherens / N. J. Conard / K. Harvati / V. Moiseyev / D. G. Drucker / J. Svoboda / M. P. Richards / D. Caramelli / R. Pinhasi / J. Kelso / N. Patterson / J. Krause / S. Pääbo / D. Reich, The genetic history of Ice Age Europe. *Nature* 534, 2016, 200-205.
- Grimm 2019: S. B. Grimm, Resilience and reorganisation of social systems during the Weichselian Lateglacial in North-West Europe. An evaluation of the archaeological, climatic, and environmental record. *Monographien des RGZM* 128 (Mainz 2019).
- Mevel/Grimm 2019: L. Mevel / S. B. Grimm, From one camp to another. First results of a comparative techno-economic analysis of the Federmesser-Gruppen lithic industries from the Central Rhineland. In: B. V. Eriksen / E. Rensink / S. Harris (eds), *The Final Palaeolithic of Northern Eurasia. Proceedings of the Amersfoort, Schleswig and Burgos UISPP commission meetings. Schriften des Museums für Archäologie Schloss Gottorf, Ergänzungsreihe* 13 (Kiel 2019) 105-135.
- Posth et al. 2016: C. Posth / G. Renaud / A. Mittnik / D. G. Drucker / H. Rougier / C. Cupillard / F. Valentin / C. Thevenet / A. Furtwängler / C. Wißing / M. Francken / M. Malina / M. Bolus / M. Lari / E. Gigli / G. Capecchi / I. Crevecoeur / C. Beauval / D. Flas / M. Germonpré / J. van der Plicht / R. Cottiaux / B. Gély / A. Ronchitelli / K. Wehrberger / D. Grigorescu / J. Svoboda / P. Semal / D. Caramelli / H. Bocherens / K. Harvati / Nicholas J. Conard / W. Haak / A. Powell / J. Krause, Pleistocene mitochondrial genomes suggest a single major dispersal of Non-Africans and a Late Glacial population turnover in Europe. *Current Biology* 26, 2016, 1-7.
- Sauer/Riede 2019: F. Sauer / F. Riede, A critical reassessment of cultural taxonomies in the Central European Late Palaeolithic. *Journal of Archaeological Method and Theory* 26, 2019, 155-184.
- Sobkowiak-Tabaka 2017: I. Sobkowiak-Tabaka, *Rozwój społeczności Federmesser na Nizinie Środkowoeuropejskiej* (Poznań 2017).
- Terberger/Barton/Street 2009: Th. Terberger / N. Barton / M. Street, The late glacial reconsidered – recent progress and interpretations. In: M. Street / N. Barton / Th. Terberger (eds), *Humans, environment and chronology of the Late Glacial of the North European Plain. RGZM – Tagungen* 6 (Mainz 2009) 189-207.
- Vaquero/Romagnoli 2018: M. Vaquero / F. Romagnoli, Searching for lazy people: The significance of expedient behavior in the interpretation of Paleolithic assemblages. *Journal of Archaeological Method and Theory* 25, 2018, 334-367.

THE ENVIRONMENTAL SETTING FOR THE LATEGLACIAL RECOLONISATION OF THE SCHELDT BASIN (NORTH-WEST BELGIUM) BY THE FEDERMESSER-GRUPPEN

The area encompassing the valleys of the River Scheldt and its tributaries, situated in North-West Belgium, is very rich in both archaeological and palaeo-environmental archives. Intensive surveys of the past decades have revealed the presence of numerous Lateglacial sites, mostly belonging to the Federmesser-Gruppen (FMG), as well as deep continuous organic soil sequences, from a variety of contexts (dune slacks, palaeo-lakes, river and stream valleys; Crombé 2006), allowing a first analysis of human behaviour against the background of a changing landscape and climate, in particular during the Allerød and the transition towards the Younger Dryas.

GENERAL SETTING

After the River Meuse, the Scheldt is the largest river in Belgium with a total length of 430 km, of which 207 km run on Belgian soils (**fig. 1**). Its headwaters are situated in northern France, its debouchment is located in the south-west of the Netherlands, where it nowadays flows into the Westerscheldt. During the Lateglacial, however, the Scheldt north of Antwerp had a more northern course, joining the estuary of the Rivers Rhine and Meuse in the central western Netherlands. The Scheldt is fed by numerous tributaries, the most important ones being from south to north Rivers Lys, Kale/Durme, Dendre and Rupel. The total catchment area amounts to 21,863 km², subdivided into the Upper Scheldt Basin (from source to Ghent) and the Lower Scheldt Basin (from Ghent to its debouchment).

The area of the Upper Scheldt, called the »Flemish Ardennes«, is a hilly upland, consisting mainly of Tertiary hills with a maximum height of 157 m above present sea-level. The quaternary cover mainly consists of loam and sandy loam deposited during the Pleniglacial, forming thick packets in the valley bottoms and on ancient river terraces. On the hill tops on the other hand the Pleistocene cover is generally thin (< 1 m), partly due to erosion, allowing Tertiary sediments to outcrop. The topography along the northern Lower Scheldt Basin is much less pronounced. This area corresponds to a typical lowland area with numerous relatively small and elongated sand ridges formed by a local reworking of coversands mainly during the late Pleniglacial and Lateglacial cold phases (Heyse 1979). In the northern and western extremes, in the Scheldt Polders and Coastal Polders, respectively, the Pleistocene coversand landscape is covered by Holocene peat and (peri)marine deposits, protecting the prehistoric sites from erosion and destruction (Crombé 2005; 2006).




Fig. 1 The River Scheldt and its tributaries with indication of the FMG sites: **1** Verrebroek »Dok 2«. – **2** Doel »Deurganckdok-sector B«. – **3** Harelbeke »Gavermeersen«. – **4** Klein-Sinaai »Boudelo«. The small box (dashed lines) indicates the position of **fig. 4**. – (Map J. Verhegge).

MATERIALS AND METHODS

Palaeo-environmental dataset

The valley of the River Scheldt, in particular its lower course, and some of its tributaries, especially the River Kale/Durme, have been studied intensively in the last decades in the context of both academic and developer-led («commercial») research. Different aspects of the Lateglacial and Early Holocene palaeo-landscapes have been intensively investigated, allowing a rather detailed reconstruction of the palaeo-vegetation through the study of pollen and plant macro-remains (Bos et al. 2017; 2018a; Deforce et al. 2005; Deforce 2011; Perdaen et al. 2011a; Storme et al. 2017; Verbruggen 1971; Verbruggen/Denys/Kiden 1996) as well as the geomorphology and palaeo-hydrology (Bogemans et al. 2012; De Moor 1963; Heyse 1979; Kiden 1989; 1991; Tavernier/De Moor 1974; Vanmaercke-Gottigny 1964). In addition, the chronological framework of the Lateglacial and Early Holocene landscape evolution is well documented by means of numerous radiocarbon dates (Crombé et al. 2012; Crombé/Robinson/Van Strydonck 2014; Meylemans et al. 2013) and a limited number of OSL dates from aeolian sediments (Bogemans/Vandenberghe 2011; Derese et al. 2010).

Archaeological dataset

The Scheldt Basin has been the subject of intense archaeological research, including surveys (field-walking, aerial photography, augering, test-pitting) and excavations, some of which covering large surfaces (e.g. Doel »Deurganckdok«, Kerkhove »Stuw«, Oudenaarde »Donk«, Verrebroek »Dok«). However, the research intensity varies considerably among the different subregions of the Scheldt Basin, resulting in somewhat biased distribution maps. Research into the Lateglacial (Final Palaeolithic) and Early Holocene (Mesolithic) archaeology has been most intense in the coversand lowland corresponding to the Lower Scheldt Basin. Systematic field walking in large parts of this area – conducted mainly by amateur archaeologists from the 1980s onwards (Crombé et al. 2011) – has led to the discovery of numerous, albeit mostly destroyed (ploughed) sites located on dry coversand ridges. The survey of the river floodplains, on the other hand, has only started in the late 1990s (Bats 2007; Bats/Bastiaens/Crombé 2006; Crombé 2006; Meylemans et al. 2013), yielding the first covered prehistoric sites in the valley of the River Scheldt and its tributaries. Some of these have been excavated in the framework of large infrastructural works, e.g. harbour expansion (Crombé 2005), water management projects (Meylemans et al. 2013; Perdaen et al. 2011b), and sand extraction (Parent/Van der Plaetsen/Vanmoerkerke 1986/1987). Lateglacial and Early Holocene archaeological research in the southern upland started already in the late 19th century, but focused almost exclusively on dryland locations, with a special interest in the hill tops and plateaus (Crombé 1989). Investigation of the valley bottoms has only been initiated recently and hence remained so far limited to occasional projects, e.g. at Spiere, Kerkhove, Oudenaarde, and Ename.

RESULTS

Late Glacial environment: Palaeo-hydrology, palaeo-topography and palaeo-vegetation

River floodplains

Within the floodplain of the Scheldt and its tributaries, remains of abandoned single-channels of meandering river systems, locally forming large oxbows, have been detected during surveys, some of which have been analysed in detail (figs 2-3). In the Scheldt these palaeo-channels locally reach dimensions of 200m width and 6-9m depth (Bogemans et al. 2012; Kiden 1989; 1991); in the Kale/Durme tributary channels of c. 30-50m width and 4-6m depth have been reported (Crombé et al. 2013; De Smedt et al. 2012). The size of these meandering channels clearly indicates that the Lateglacial rivers had a discharge at least three to five times larger than today (Kiden 1989; 1991). This is also confirmed by the dimensions of the Lateglacial palaeo-meanders, which were much larger compared to present-day oxbows, locally extending 3-4.5km inland. The higher discharge during the Lateglacial probably is due to higher levels of meltwater in spring and a less dense vegetation which induced more intense run-off (Kiden 1991).

The exact dating of the initial incision of these meandering channel systems is still not well established. The basal infill of sandy gyttja or scroll-bar sediments in two deep channels – one at Kalken in the Lower Scheldt Valley (Meylemans et al. 2013) and another at Vinderhoute in the Upper Kale Valley (Verbruggen 1971) – yielded ages of $12,460 \pm 60$ years ¹⁴C-BP (Beta-245745) and $12,655 \pm 70$ years ¹⁴C-BP (GrN-6062), respectively. These situate the start of the infilling during the Bølling (GI-1e) or even slightly earlier. However, given the presence of numerous reworked pollen, e.g. from several thermophilous (*Corylus*, *Carpinus*) and Tertiary species, as well as the high percentages of *Pinus*, the reliability of these early radiocarbon dates is





Fig. 2 Above: Lateglacial palaeo-meanders along the Lower Scheldt and Lower Lys Rivers: **A** palaeo-meanders with fossil scroll-bars; **B** river dunes; **C** present-day rivers. – Left: detail of the Lateglacial palaeo-meander of Overmere-Donk: **1** border of the Lateglacial floodplain; **2** fossil Lateglacial river channel; **3** scroll-bars; **4** present-day river bed; **5** river dunes; **6** altitude in m TAW (= mean low water tide level in Oostende). – (After Kiden/Verbruggen 2001, fig. 7).

generally contested (Kiden 1991; Storme et al. 2017; Verbruggen 1971), as they may have been performed on eroded older material.

The first reliable radiocarbon dates all come from calcareous gyttja found in several channels immediately above the sandy basal sediments. In the Lower Scheldt Valley the oldest dates are around $11,120 \pm 60$ years ^{14}C -BP (Beta 245744) and $10,910 \pm 60$ years ^{14}C -BP (Beta 245743; Meylemans et al. 2013), corresponding to the final Allerød (GI-1a) and the beginning of the Younger Dryas (GS-1). Similar dates have been obtained from several channel sections in the Kale/Durme Valley (Bos et al. 2018b; Crombé et al. 2013; Crombé/Robinson/Van Strydonck 2014). With the exception of one date from Ename ($11,210 \pm 50$ years ^{14}C -BP; Deforce 2004), no radiocarbon dates are yet available for the Upper Scheldt Valley, but pollen evidence




Fig. 3 Schematic cross-sections of the valleys of the Upper Scheldt (A) and Lower Scheldt (B): **1** braided river channel-fill and bar deposits. – **2** fining upward overbank deposits. – **3** aeolian deposits. – **4** Lateglacial meandering river lateral accretion deposits forming scroll-bars. – **5** Subatlantic meandering river lateral accretion deposits. – **6** gyttja. – **7** peat and clayey peat. – **8** fluvatile clay and silt. – **9** estuarine intertidal deposits. – **10** man-made embankments. – (After Kiden 1991, fig. 12.3).

points to an initial infilling from the start of the Younger Dryas (Parent/Van der Plaetsen/Vanmoerkerke 1986/1987). So although the timing of the incision of the meandering channels is not yet precisely fixed, it is clear that it occurred before the end of the Allerød/start of the Younger Dryas when the final channels started to fill in locally. This is in agreement with data from other river valleys, such as the Somme (Antoine et al. 2000), the Meuse (van Huissteden/Kasse 2001), and the Niers-Rhine Valley (Kasse et al. 2005). In all these valleys the incision of a large single channel meandering system has been dated to the Allerød, while the Bølling is characterised as a transitional phase between a braided and a meandering system. Remains of such a transitional system, characterised by several straight, small, and shallow channels which were active at the same time, have only been found so far within the Moervaart depression along the Kale/Durme River (Crombé et al. 2013; De Smedt et al. 2012) (fig. 4). Radiocarbon dates from a series of these anastomosing channels demonstrate that the infilling started at the transition from the GS-2a (Oldest Dryas) to the GI-1e (Bølling) or early in the GI-1e between $12,450 \pm 50$ years ^{14}C -BP (Beta-302750) and $11,925 \pm 55$ years ^{14}C -BP (KIA-47009), and continued during the main part of the Allerød (GI-1c) (Crombé/Robinson/Van Strydonck 2014). However, some channels may be younger as indicated by a late Allerød date from the basis of a small anastomosing channel at Wachtebeke ($11,345 \pm 50$ years ^{14}C -BP; KIA-46184; Crombé et al. 2013; Crombé/Robinson/Van Strydonck 2014).

The later evolution of the Scheldt River system, however, deviates considerably from the river valleys in surrounding countries. In the case of the Meuse the cold conditions of the Younger Dryas led to a return to a multi-channel braided system followed by a new deep channel incision at the start of the Holocene (Hoek/Bohncke 2002), while in the case of the Somme the Younger Dryas coincides with a major infilling of the alluvial plain followed also by a new major channel incision at the start of the Preboreal. On the contrary, in the Scheldt Basin the Allerød meandering channel system remained active during the Younger Dryas and




Fig. 4 Map of the Moervaart palaeo-lake through which the single-channel meandering River Kale/Durme runs (underlying map = digital elevation model; Lidar, AGIV 2001-2004). The dots indicate the position of FMG sites. In the west, there are remains of an older, anastomosing river system consisting of several small and shallow channels, reconstructed on the basis of data from augering, electromagnetic interference (EMI) and digital elevation modelling. Most FMG sites along these shallow channels are situated on small levees. In contrast, no sites are known along the lower Kale/Durme River in the east.

into the Holocene (Bogemans et al. 2012; Crombé et al. 2013; Crombé/Robinson/Van Strydonck 2014; Kiden 1991; Storme et al. 2017); however, with a clear change in sedimentation from calcareous gyttja to peat in the course of the Preboreal. The latter was probably the result of a lowering of the water level caused by increased evapotranspiration following the installation of a dense pine (*Pinus*) forest as well as higher temperatures. The sedimentary characteristics indicate an evolution from deep, stagnant, or slow-running water during the late Allerød and early parts of the Younger Dryas to very shallow and marshy conditions during the later Younger Dryas and the beginning of the Holocene. Due to the reduced fluvial activity during the Younger Dryas, river (parabolic) dunes were created within or just outside the Lateglacial floodplain. Some of these were OSL dated to $12,000 \pm 900$ years (OSL)-BP ($n=5$) (Bogemans/Vandenberghe 2011).

Sandy lowland (Lower Scheldt Basin)

The landscape in the northern sandy lowland was very dynamic as a result of a changing climate during the Lateglacial. At the start, local reworking of Pleniglacial (cover)sands triggered the formation of dunes, a process which was reactivated during the subsequent cold Dryas stages (Heyse 1983). Most dunes are rather small, except for one called »the Great Sand Ridge of Maldegem-Stekene«, running over a distance of c. 80km from west to east across the sandy lowland (Heyse 1979; 1983). According to recent data (Bos et al. 2013; Crombé et al. 2012), this massive dune was formed mainly during the Older Dryas (GI-1d) and Younger Dryas (GS-1) and consists of numerous intersecting and overlapping smaller dunes. In between these

sand dunes hundreds of relatively small- to medium-sized, closed, and shallow depressions were formed by local aeolian erosion, so-called blow-outs. Judging by the presence of humiferous to peaty sediments at their base, it can be concluded that these shallow depressions were temporarily wet, ranging from ponds during the Bølling to shallow dune slacks in the Allerød (Bos et al. 2013; Crombé et al. 2012).

Another type of open-water system within the sandy lowland consisted of shallow freshwater lakes which appeared along the southern steep edge of »the Great Sand Ridge«. Contrary to the dune slacks, these were not formed by wind erosion, but by local accumulation of groundwater and surface water which had become blocked by »the Great Sand Ridge« (De Moor/Heyse 1978; Verbruggen/Denys/Kiden 1996). By far the largest one was Lake Moervaart extending over a surface of approximately 25 km² (Heyse 1983; Crombé et al. 2013) (fig. 4). Starting as swampy depressions during the Bølling, they evolved in the course of the Allerød into shallow lakes with fluctuating water levels (3-6 m), probably as a result of increased precipitation and local seepage (Bos et al. 2013; Crombé et al. 2013; Denys/Kiden/Verbruggen 1998). The highest water level in the Moervaart Lake was reached at the transition from the early to the middle Allerød, as indicated by the abundance of aquatic plant species, such as waterlily (*Nymphaea*), Eurasian watermilfoil (*Myriophyllum*), and buckbean (*Menyanthes trifoliata*) (Bos et al. 2017).

The vegetation around these dune slacks and lakes gradually became more closed in the course of the Allerød (Bos et al. 2018a). Initially birch (*Betula*) and willow (*Salix*) were the dominant tree species, although grasses and sedges remained largely present along the banks. From c. 11,300 years ¹⁴C-BP onwards Scots pine (*Pinus sylvestris*) increased in importance and finally became the most frequent species. By the end of the Allerød the Arboreal Pollen ratio locally reached 80-90 %, indicating the presence of an almost entirely closed coniferous forest (Bos et al. 2013; 2018a; Deforce et al. 2005; Verbruggen/Denys/Kiden 1996).

Clearly before the onset of the colder Younger Dryas, more precisely during the *Pinus*-stage of the Allerød, a clear decrease of the water level occurred both in the dune slacks and lakes. Dune slacks got filled in with aeolian sands (Bos et al. 2013; Crombé et al. 2012), while in the lakes (Lakes Moervaart and Snellegem) the lacustrine sediments got covered by peat indicating a change to a marshier environment (Denys/Verbruggen/Kiden 1990; Denys/Kiden/Verbruggen 1998). This marked lowering of the water level was recently synchronised with the Intra-Allerød Cold Period (GI-1b) on the basis of a Bayesian modelling of radiocarbon dates (Crombé/Robinson/Van Strydonck 2014), yet the chronological precision still demands further improvement.

Loamy upland (Upper Scheldt Basin)

From the southern upland area no palaeo-environmental information is currently available, except for the Scheldt floodplain (see above). This is mainly due to erosion of Lateglacial levels situated on hill tops and slopes, as well as plateau edges. In addition, so far only little environmental research has been conducted in small stream valleys (Maarkebeek, Zwalmbeek, etc.) that feed the River Scheldt and its tributaries.

FMG occupation

Lithic assemblages

Except for some isolated late Magdalenian/Creswellian finds in the upland area (Crombé 1989; Vandendriessche et al. 2016), there are currently no traces of human occupation in the Scheldt Basin prior to the




Fig. 5 Selection of backed points from the Scheldt Basin attributed to different stages of the Allerød and Younger Dryas: **1-3** early FMG (Doel »Deurganckdok«). – **4-5** late FMG (Stekene »Heirweg«). – **6-9** final FMG (Malaure-like point: Klein-Sinaai), and Younger Dryas traditions (Blanchères point: Evergem »De Nest«; microlithic armatures: Ruien »Rosalinde«). – (Drawings J. Sergant).

FMG. A clear explanation for this absence of human occupation during the Bølling has not yet been found, but it could be that the environment was still too unstable for humans to settle. The sand dunes were still in formation and probably not yet well-developed, making the area not well suited for erecting camp sites. In addition, there may have been a problem with the availability of drinking water, certainly in the sandy lowland where the dune slacks and lakes only appeared later during the Allerød (fig. 5).

At the moment it looks as if the FMG hunter-gatherers were the first re-colonisers of the Scheldt Basin after the Lateglacial Maximum. At least 30 FMG sites are actually known in the Scheldt Basin (Ameels/Van Vlaenderen 1995; Crombé/Verbruggen 2002; Crombé et al. 2011) (fig. 1). Except for a few, these are all eroded plough-layer sites known from field surveys. As a result little is currently known about the chronology and settlement organisation of the FMG in the Scheldt Basin. The lithic assemblages found on these surface sites are generally limited in size, ranging between 500 and 1,000 artefacts. The main raw material consists of local flint of mediocre quality found as relatively small nodules on Tertiary outcrops. The use of better quality, black fine-grained flint of Obourg-type is generally restricted to less than 10 % (Crombé et al. 2011). Besides backed points and blade(let)s the lithic toolkit comprises standardly numerous scrapers and burins, the latter mainly dihedral and made on truncation.

So far just four sites have yielded *in situ* finds. The site of Verrebroek »Dok 2« (Crombé et al. 1999; Crombé 2005) consists of a very small knapping post (< 1 m²) without diagnostic tools, found along a former dune slack on top of a humiferous layer dated by pollen analysis and radiocarbon method to the Older Dryas and/or the start of the Allerød (Crombé et al. 2012; Van Strydonck 2005). A radiocarbon date obtained on a sample of burnt bark found in the same stratigraphical position as the flint artefacts yielded a result of 11,900 ± 90 years ¹⁴C-BP (UtC-9434), corresponding to the Older Dryas/Allerød transition (Van Strydonck/Crombé 2005). This date is also corroborated by the knapping characteristics which indicate careful and rather intense core preparation (frequent rejuvenations, high frequency of faceted butts, etc.) and the use of a soft stone hammer (Perdaen/Crombé/Sergant 2004; Perdaen/Ryssaert 2002). These features are typical of early rather than late FMG in Western Europe (Bodu/Valentin 1997; Fagnart 1997). Pollen and charcoal analyses (Deforce et al. 2005) indicate that the occupation took place in an environment with up to 45 % of willow (*Salix*) and a wet vegetation consisting mainly of *Cyperaceae* (sedge family) and *Poaceae* (grass family).

At the site of Doel »Deurganckdok-sector B« (Crombé et al. 2000; Crombé 2005) FMG lithic remains have been found on a coversand ridge covered by Holocene peaty and fluvial deposits. Despite this covering, large parts of the site were disturbed by intense later occupations dated to the Late Mesolithic (> 100 hearth-pits) and Early Neolithic (Swifterbant Culture). In addition – due to advanced bioturbation – the FMG remains were intermixed with lithics from these younger phases. However, based on a combination of attributes, such as raw material, weathering (patination), and morphology, it was possible to select c. 600 artefacts which can be attributed to the FMG with a high degree of certainty. Despite the absence of radiocarbon dates, here too, the knapping characteristics and the presence of some »bipointes« (fig. 5, 1-3) seem to point to an early stage of the FMG, although the presence of late FMG artefacts is not excluded (Perdaen/Crombé/Sergant 2004).

A third excavated site is situated within the extensive site-complex along the northern bank of the Moervaart palaeo-lake (see below). At Klein-Sinaai »Boudelo« (Vanmoerkerke/De Belie 1984) at least five lithic scatters were discovered by chance during the excavation of a Medieval abbey. According to a first preliminary excavation report, the FMG remains (1,757 lithic artefacts) were found in a sandy layer with iron nodules, situated c. 10 cm above a layer containing scattered charcoal fragments. The latter might correspond to the well-known Usselo-layer, but this unfortunately cannot be verified since the site is not accessible any longer and the charcoal has not been preserved. No further spatial information is available, but in all scatters some typical FMG tools (backed points and blades, burins) have been reported. Worth mentioning are two »bipointes« and one »Mauricie-like point« (fig. 5, 6), which might point to an early and final FMG occupation of the site, respectively.

A last FMG site was excavated at Harelbeke »Gavermeersen« in the context of sand extraction (Vermeersch 1976). Here FMG hunter-gatherers choose to settle on the dry northern bank of a large depression, situated

less than 1.5 km from the Lys floodplain, an important tributary of the River Scheldt. This shallow depression was filled with silty to clayey sediments sealed by peat, which unfortunately has neither been studied nor dated. The site itself (c. 2,183 lithic artefacts) is situated on the southern flank of a sandy ridge, which was largely affected by ploughing. Nevertheless, some lithics were still preserved underneath the plough-layer, albeit not allowing a meaningful spatial analysis.

Although the above excavations have revealed the presence of the early stages of FMG, in terms of overall technology the vast majority of lithic assemblages in the Scheldt Basin clearly relates to the classic/late FMG, characterised by uni-pointed backed armatures (**fig. 5, 4-6**) and a much less refined blade technology that focused on the production of irregular unstandardised blade(lets).

Distribution pattern

The distribution pattern of the FMG sites displays a marked concentration in the sandy lowland area (**fig. 1**). In the southern uplands FMG sites are currently hardly known, but this may be partly due to a lesser research intensity and taphonomic factors. Indeed, one needs to consider the possibility that numerous sites in this hilly area are hidden underneath colluvial deposits, as was recently attested at the Younger Dryas site of Ruien »Rosalinde« (Crombé et al. 2014). The few sites known so far in the uplands (e.g. Kluisberg, Pottelberg, Kemmelberg) are located on hill tops, in particular on local outcrops of Tertiary sands (Vandendriessche et al. 2016). A similar observation has been reported by Coudret and Fagnart (2006, 737) for the Somme region of northern France.

Within the sandy lowland region of the Lower Scheldt Basin FMG sites mainly occur along the dry banks of dune slacks and shallow lakes, sometimes forming extensive site-complexes running over several kilometres (Crombé et al. 2011; 2013). This pattern is most apparent along the Moervaart palaeo-lake (**fig. 4**), where a high density of sites has been observed along the relatively steep northern bank and on small sandy ridges, possible levees, within the western lake sector. These dense clusters of sites probably reflect frequent and recurrent visits of FMG hunter-gatherers to these lake areas. Most likely they were attracted by the rich and diversified ecology of these areas, which provided them with drinking water, rich hunting grounds, and plenty of plant material for both consumption and tool production (Crombé et al. 2013).

In contrast to this, the total absence of FMG sites along the floodplains of the River Scheldt and its tributaries is most remarkable and difficult to explain. Despite the intense archaeological research especially along the Lower Scheldt Valley, so far not a single FMG camp-site, except some isolated finds, has been discovered within the Lateglacial floodplains. Apparently the scroll-bars flanking the Allerød palaeo-channels were inhabited only from the Early (Boreal) Holocene onwards, when the rivers were already reduced to shallow slow-running streams (Ameels et al. 2003; Bats 2005; 2007; Bats/Bastiaens/Crombé 2006; Crombé et al. 2013; Meylemans et al. 2013; Parent/Van der Plaetsen/Vanmoerkerke 1986/1987; Perdaen et al. 2011a; 2011b). Even on the dry river dunes situated adjacent to the floodplains not one FMG site is currently known, while the evidence of Mesolithic and Neolithic occupation is overwhelming. This is particularly evident along the very intensively surveyed Durme River (Crombé et al. 2011; 2013; **fig. 4**, eastern section). Here field-walking yielded at least twelve Early (Boreal) Mesolithic sites, mostly situated on the left bank, while not a single FMG site, not even an isolated find, has been reported. Similarly, intensive surveys and large-scale excavations on the dry banks along the floodplains along the Upper Scheldt, e.g. at Ename (Ameels et al. 2003), Kerkhove (Crombé 1985; Sergeant et al. 2016), Spiere (Vanmoerkerke 1988; Vanmontfort et al. 2001/2002), Gavere (Vanmoerkerke 1986) and Melden (Crombé/Braeckman/Parent 1991), yielded plenty of Mesolithic and extensive Neolithic (Michelsberg Culture) sites, but no evidence of FMG occupation.

DISCUSSION

From the above it is clear that the Lateglacial environment in the Scheldt Basin was very dynamic. At present not all details of the Lateglacial environmental changes are fully understood, but the main trends are yet defined. During the time of occupation of the FMG, i.e. the Allerød, an important network of single-channel meandering rivers, with the Scheldt as the main river, as well as numerous shallow ponds and lakes had developed within an increasingly wooded environment dominated by first birch and later pine. Within this landscape FMG hunter-gatherers clearly preferred to settle along the banks of lakes and ponds, apparently leaving the river valleys aside. This is a very remarkable pattern which considerably differs from surrounding river valleys in Western Europe, where FMG hunter-gatherers intensively occupied the edges of Lateglacial floodplains. In the Dutch Meuse Valley FMG sites occur both along lake banks and the outer banks of Lateglacial oxbows (Deeben 1995). In northern France, the Somme and Seine Valleys have yielded plenty of sealed FMG sites situated along active river channels (Bodu/Debout/Bignon 2006; Coudret/Fagnart 2006). Similarly in the valley of the Rhine and its tributaries FMG sites, such as Andernach, Bad Breisig, Niederbieber and Kettig, are located close to active meandering channels (Street et al. 2006).

Why do the River Scheldt and its tributaries differ from this general pattern? Before addressing this question it needs to be investigated whether the actual distribution pattern is not biased. It has to be considered that some FMG sites may be hidden as a result of later sedimentation. As a matter of fact, most sites in adjacent valleys are covered by later sediments of different origin (alluvial, volcanic, aeolian, anthropogenic). However, it seems highly unlikely that this explains the total lack of FMG sites in the Scheldt floodplains. The small scroll-bars created as a result of the lateral migration of the meandering channels, which constituted the preferred habitats for human occupation during the Mesolithic, are often hardly covered, some are even visible in the present day landscape. So, if FMG sites are present on these small elevations, at least some of them would have been found during the intense surveys of the Scheldt floodplains. FMG remains were also not reported on the numerous drilled and/or excavated scroll-bars which are covered by Holocene fluvial deposits, e.g. at Oudenaarde, Kerkhove, Kalken, Wichelen, Bazel, and Melsele. Still, it cannot be fully excluded that some FMG sites have been missed during archaeological research. The discovery of a humiferous palaeo-sol situated at maximum 30 cm below the top of a scroll-bar at Wichelen »Bergenmeersen« (Perdaen/Meylemans/Vanholme 2013) demonstrates that some scroll-bars might have been covered by aeolian sands during the Younger Dryas. Knowing that archaeological research is often limited to the top of the Pleistocene sediments, it is possible that some covered FMG levels were not explored, but this certainly cannot explain the complete absence of sites in the floodplains. Aeolian sedimentation could also be potentially responsible for the lack of FMG sites on the dry banks along the margins of the floodplains. However, if the river dunes along the Scheldt and its tributaries were created during the Younger Dryas, as suggested by recent OSL research on some of them (Bogemans/Vandenberghe 2011), they did not yet exist at the time of the FMG. So, at present it rather looks as if the banks of the Lateglacial floodplains and oxbows were not yet attractive for settling during the Allerød due to the absence of dry enough settlement locations, comparable to the inland dunes intensively occupied by FMG groups. Some isolated finds of FMG armatures nevertheless indicate that the Scheldt floodplains were visited by FMG hunter-gatherers, possibly in the context of hunting activities.

Alternatively we need also to consider post-Allerød fluvial erosion of former FMG sites in order to explain the lack of sites in the floodplains. However, this explanation too seems rather unlikely as, contrary to other valleys in North-West Europe, the Scheldt Valley remained rather stable and much less dynamic during the subsequent Younger Dryas and Early Holocene. The single-channel systems from the Allerød persisted into the Holocene with a gradual infilling of the deep channels demonstrating a marked decrease of the fluvial

activity. In fact the Younger Dryas and Early Holocene rivers were largely reduced to slow running streams. It is only during the early stages of the Subatlantic that a new river channel with associated crevasse gullies was incised in the Scheldt floodplain (Bogemans et al. 2012; Kiden 1991). However, sedimentary properties seem to indicate a rather stable meandering channel with minor lateral migration, so the impact of this renewed fluvial activity on older prehistoric sites will have been rather limited.

So, actually there is little evidence which supports the idea that the distribution pattern of FMG sites at least along the Lower Scheldt Basin is seriously biased. Then how can we explain the absence of interest in the river floodplains? Maybe the river valleys were a too dynamic environment for FMG hunter-gatherers in order to exploit or settle? Large parts of the Allerød coincide with a phase of high discharge and fluvial erosion; river channels were not yet fixed but subject to continuous lateral migration, which ultimately led to the formation of very extensive oxbows. However, this argument does not hold for the later phases of the Allerød. The available radiocarbon evidence clearly indicates that the river systems in the Scheldt Basin stabilised ultimately during the final Allerød (GI-1a) and the transition towards the Younger Dryas (GS-1). At that time the channels started to fill in with organic sediments, indicating more tranquil flow rates. So one would at least expect FMG hunter-gatherer activity along the Scheldt and its tributaries at this late stage, even more so since around the same time the coversand interior faced an abrupt drop of the ground water level, turning most of the inland freshwater lakes and dune slacks into dry or marshy environments. This certainly must have had a considerable impact on the FMG hunter-gatherers who intensively exploited these lake banks. They were confronted with a sudden reduction of the available open water, which provided them with vital drinking water. Also the game populations must have been affected by this abrupt environmental change. The frequent occurrence of animal dung (coprophilous) fungi (*Bombardioidea* type, *Sporormiella* type, *Podospora* type, *Sordaria* type and other Sordariaceae; Bos et al. 2013) as well as high phosphate ratios (Louwagie/Langohr 2005, 103) indirectly demonstrate the presence of numerous large herbivorous mammals along the Allerød lake banks. These too will have suffered from the marked reduction in available drinking water and will have been forced to move to other areas. In this context the total absence of FMG sites along the main river valleys of the Scheldt Basin is very strange. Does it mean that these hunter-gatherers moved to other areas, altogether leaving the Scheldt Basin? Unfortunately, due to the complete absence of radiocarbon dates for the FMG, this hypothesis cannot yet be further explored but nevertheless seems plausible. It seems reasonable to assume that the important environmental changes occurring during the final Allerød and early Younger Dryas altered the settlement system of the contemporaneous hunter-gatherers, but further research is needed in order to fully understand the nature of the human response.

CONCLUSIONS

The most important observation from the above analysis is the so far total absence of FMG camp sites along the river floodplains of the Scheldt Basin. Apparently the first re-colonisers of North-West Belgium were more attracted to the dry banks of former freshwater lakes and ponds in the sandy interior, possibly indicating that the latter constituted a more stable and secure environment, contrary to the highly dynamic river valleys – at least until the final Allerød (GI-1a) when a sudden ground water lowering dramatically reduced the available amount of drinking water for both animals and humans. But even then, FMG hunter-gatherers apparently did not shift to the rivers, although these probably had become the only or at least the most important sources of drinking water within the area. However, in absence of faunal remains and well-dated FMG sites it remains difficult to assess the real extent of this hydrological event. Future Lateglacial research in the Scheldt Basin should therefore focus on the detection and excavation of sealed FMG sites, which can

still be expected along the banks of the numerous freshwater lakes and dune slacks. Sites such as Verrebroek »Dok 2« and Klein-Sinaai »Boudelo« clearly prove the existence of well-preserved FMG occupation levels underneath Younger Dryas aeolian sediments. Existing survey techniques should be further refined, allowing to map buried palaeo-landscapes on a larger scale and with a higher resolution (Missiaen et al. 2015; Verhegge/Missiaen/Crombé 2016) and to detect even the smaller and less dense sites (Crombé/Verhegge 2015). In addition much more research should be done on the contemporaneous environment in view of a better understanding of the landscape evolution, especially in response to a changing climate. The long and often continuous sedimentary sequences available in the numerous river channels and inland lakes make the Scheldt Basin an ideal context for a high-resolution, multi-proxy investigation of the Lateglacial and Early Holocene environment.

Acknowledgements

I wish to thank Patrick Kiden for allowing me to reprint **figs 2-3**. Thanks also to Jeroen Verhegge and Joris Sergeant for producing **figs 1** and **5**, respectively. Lastly, the author also wants to acknowledge the reviewers whose feedback has contributed to improving the quality of this paper.

References

- Ameels/Van Vlaenderen 1995: V. Ameels / L. Van Vlaenderen, Epipaleolithische en vroeg-mesolithische sites langsheen de Moervaart (Oost-Vlaanderen). *Notae Praehistoricae* 15, 1995, 35-43.
- Ameels et al. 2003: V. Ameels / J. Bastiaens / M. Bats / Ph. Crombé / K. Deforce / K. Haneca / J.-P. Parent / M. Van Strydonck, Recent Steentijdonderzoek in de regio Oudenaarde (Oost-Vlaanderen, België). *Notae Praehistoricae* 23, 2003, 61-65.
- Antoine et al. 2000: P. Antoine / J.-P. Fagnart / N. Limondin-Lozouet / A. V. Munaut, Le Tardiglaciaire du bassin de la Somme: éléments de synthèse et nouvelles données. *Quaternaire* 11, 2000, 85-98.
- Bats 2005: M. Bats, Prospectief booronderzoek in de Kalkense Meersen (prov. Oost-Vlaanderen, België). *Notae Praehistoricae* 25, 2005, 203-207.
- 2007: M. Bats, The Flemish Wetlands: an archaeological survey of the valley of the river Scheldt. In: J. Barber / C. Clark / M. Cressey / A. Crone / A. Hale / J. Henderson / R. Housley / R. Sands / A. Sheridan (eds), *Archaeology from the Wetlands: Recent Perspectives. Proceedings of the 11th WARP Conference, Edinburgh 2005. Society of Antiquaries of Scotland. WARP Occasional Paper 18 (Edinburgh 2007)* 93-100.
- Bats/Bastiaens/Crombé 2006: M. Bats / J. Bastiaens / Ph. Crombé, Prospectie en waardering van alluviale gebieden langs de Boven-Schelde, CAI-project 2003-2004. In: E. Meylemans (ed.), *VIOE-rapporten 02, Centrale Archeologische Inventaris CAI-II, »Thematisch inventarisatie- en evaluatieonderzoek«* (Brussel 2006) 75-100.
- Bodu/Valentin 1997: P. Bodu / B. Valentin, Groupes à Federmesser ou Aziliens dans le sud et l'ouest du bassin parisien. Propositions pour un nouveau modèle d'évolution. *Bulletin de la Société préhistorique française* 94, 1997, 341-347.
- Bodu/Debout/Bignon 2006: P. Bodu / G. Debout / O. Bignon, Variabilité des habitudes tardiglaciaires dans le Bassin parisien: l'organisation spatiale et sociale de l'Azilien ancien du Clouseau. *Bulletin de la Société préhistorique française* 103, 2006, 711-728.
- Bogemans/Vandenberghe 2011: F. Bogemans / D. Vandenberghe, OSL dating of an inland dune along the lower River Scheldt near Aard (East Flanders, Belgium). *Netherlands Journal of Geosciences* 90, 2011, 23-29.
- Bogemans et al. 2012: F. Bogemans / E. Meylemans / J. Jacobs / Y. Perdaen / A. Storme / I. Verdurmen / K. Deforce, The evolution of the sedimentary environment in the lower River Scheldt valley (Belgium) during the last 13,000 a BP. *Geologica Belgica* 15, 2012, 105-112.
- Bos et al. 2013: J. A. A. Bos / F. Verbruggen / S. Engels / Ph. Crombé, The influence of environmental changes on local and regional vegetation patterns at Rieme (NW Belgium): implications for Final Palaeolithic habitation. *Vegetation History and Archaeobotany* 22, 2013, 17-38.
- 2017: J. A. A. Bos / Ph. De Smedt / H. Demiddele / W. Z. Hoek / R. Langohr / V. Marcelino / N. Van Asch / D. Van Damme / Th. Van der Meeren / J. Verniers / P. Boeckx / M. Boudin / M. Court-Picon / P. Finke / V. Gelorini / S. Gobert / O. Heiri / K. Martens / F. Mostaert / L. Serbruyns / M. Van Strydonck / Ph. Crombé, Multiple oscillations during the Lateglacial as recorded in a multi-proxy, high-resolution record of the Moervaart palaeolake (NW Belgium). *Quaternary Science Reviews* 162, 2017, 26-41.
- 2018a: J. A. A. Bos / Ph. De Smedt / H. Demiddele / W. Z. Hoek / R. Langohr / V. Marcelino / N. Van Asch / D. Van Damme / Th. Van der Meeren / J. Verniers / Ph. Crombé, Weichselian Lateglacial environmental and vegetation development in the Moervaart palaeolake area (NW Belgium); implications for former human

- occupation patterns. Review of Palaeobotany and Palynology 248, 2018, 1-14.
- 2018b: J. A. A. Bos / V. Gelorini / Th. Van der Meeren / J. Peleman / M. Court-Picon / H. Demiddele / Ph. De Smedt / M. Boudin / J. Verniers / P. Boeckx / Ph. Crombé, The Younger Dryas and Preboreal landscape in the Moervaart area (northwestern Belgium) and the apparent decrease in human occupation. Vegetation History and Archaeobotany 27, 2018, 697-715.
- Coudret/Fagnart 2006: P. Coudret / J.-P. Fagnart, Données préliminaires sur les habitats des groupes de la tradition à Federmesser du bassin de la Somme. Bulletin de la Société préhistorique française 103, 2006, 729-740.
- Crombé 1985: Ph. Crombé, Een prehistorisch site te Kerkhove (Mesolithicum-Neolithicum). Westvlaamse Archaeologica 1, 1985, 3-39.
- 1989: Ph. Crombé, Découvertes pré- et protohistoriques dans »la region des collines«. Cahiers de Préhistoire du Nord 6, 1989, 15-21.
- 2005: Ph. Crombé (ed.), The last hunter-gatherer-fishermen in Sandy Flanders (NW Belgium). The Verrebroek and Doel excavation projects. 1: Palaeo-environment, Chronology and Features. Archaeological Reports Ghent University 3 (Gent 2005).
- 2006: Ph. Crombé, The wetlands of sandy Flanders (Northwest Belgium): potentials and prospects for prehistoric research and management. In: E. Rensink / H. Peeters (eds), Preserving the Early Past. Investigation, Selection and Preservation of Palaeolithic and Mesolithic Sites and Landscapes. Nederlandse Archeologische Rapporten 31 (Amersfoort 2006) 41-54.
- Crombé/Verbruggen 2002: Ph. Crombé / C. Verbruggen, The Late-glacial and early Postglacial occupation of northern Belgium: the evidence from Sandy Flanders. In: B. V. Eriksen / B. Bratlund (eds), Recent studies in the Final Palaeolithic of the European plain. Proceedings of a UISPP Symposium, Stockholm, 14.-17. October 1999. Jutland Archaeological Society Publications 39 (Højbjerg 2002) 165-180.
- Crombé/Verhegge 2015: Ph. Crombé / J. Verhegge, In search of sealed Palaeolithic and Mesolithic sites using core sampling: the impact of grid size, meshes and auger diameter on the discovery probability. Journal of Archaeological Science 53, 2015, 445-458.
- Crombé/Braeckman/Parent 1991: Ph. Crombé / K. Braeckman / J.-P. Parent, Baggervondsten afkomstig van een mesolithische en neolithische site te Melden »Waarde« (Oudenaarde, O.-VI.). Handelingen van het Zottegems Genootschap voor Geschiedenis en Oudheidkunde 5, 1991, 211-234.
- Crombé/Robinson/Van Strydonck 2014: Ph. Crombé / E. Robinson / M. Van Strydonck, Synchronizing a Late Glacial Abrupt Cooling Event with Paleoenviromental and Population Changes: Case Study of the Moervaart Paleolake Area (NW Belgium). Radiocarbon 56, 2014, 899-912.
- Crombé et al. 1999: Ph. Crombé / K. Deforce / R. Langohr / G. Louwagie / Y. Perdaen / J. Sergeant / C. Verbruggen, A small Final Palaeolithic knapping site at Verrebroek »Dok 2« (Flanders, Belgium). Notae Praehistoricae 19, 1999, 63-68.
- 2000: Ph. Crombé / J.-P. Van Roeyen / J. Sergeant / Y. Perdaen / K. Cordemans / M. Van Strydonck, Doel »Deurganckdok« (Flanders, Belgium): settlement traces from the Final Palaeolithic and the Early to Middle Neolithic. Notae Praehistoricae 20, 2000, 111-119.
- 2011: Ph. Crombé / J. Sergeant / E. Robinson / J. De Reu, Hunter-gatherer responses to environmental change during the Pleistocene-Holocene transition in the southern North Sea basin: final Palaeolithic – Final Mesolithic land use in northwest Belgium. Journal of Anthropological Archaeology 30, 2011, 454-471.
- 2012: Ph. Crombé / M. Van Strydonck / M. Boudin / T. Van den Brande / C. Derese / D. A. G. Vandenberghe / P. Van den Haute / M. Court-Picon / J. Verniers / J. A. A. Bos / F. Verbruggen / M. Antrop / M. Bats / J. Bourgeois / J. De Reu / Ph. De Maeyer / Ph. De Smedt / P. A. Finke / M. Van Meirvenne / A. Zwertvaegher, Absolute dating (¹⁴C and OSL) of the formation of coversand ridges occupied by prehistoric hunter-gatherers in NW Belgium. Radiocarbon 54, 2012, 715-726.
- 2013: Ph. Crombé / Ph. De Smedt / N. S. Davies / V. Gelorini / A. Zwertvaegher / R. Langohr / D. Van Damme / M. Van Strydonck / M. Antrop / J. Bourgeois / Ph. De Maeyer / J. De Reu / P. A. Finke / M. Van Meirvenne / J. Verniers, Hunter-gatherer responses to the changing environment of the Moervaart palaeolake (NW Belgium) during the Late Glacial and Early Holocene. Quaternary International 308/309, 2013, 162-177.
- 2014: Ph. Crombé / J. Sergeant / A. Verbrugge / A. De Graeve / B. Cherretté / J. Mikkelsen / V. Cnudde / T. De Kock / D. J. Huisman / B. J. H. van Os / M. Van Strydonck / M. Boudin, A sealed flint knapping site from the Younger Dryas in the Scheldt valley (Belgium): bridging the gap in human occupation at the Pleistocene-Holocene transition in W Europe. Journal of Archaeological Science 50, 2014, 420-439.
- Deeben 1995: J. Deeben, Human occupation of the southern Netherlands during the Younger Dryas (extended abstract). Geologie en Mijnbouw 74, 1995, 265-269.
- Deforce 2004: K. Deforce, Rapport palynologisch onderzoek van een fossiele Scheldegeul te Oudenaarde-Stuw (Oudenaarde). Internal Report Ghent University (Ghent 2004).
- 2011: K. Deforce, Middle and Late Holocene vegetation and landscape evolution of the Scheldt estuary. A palynological study of a peat deposit from Doel (N-Belgium). Geologica Belgica 14, 2011, 277-288.
- Deforce et al. 2005: K. Deforce / V. Gelorini / C. Verbruggen / L. Vrydaghs, Pollen and phytolith analyses. In: Crombé 2005, 108-126.
- De Moor 1963: G. De Moor, Bijdrage tot de kennis van de fysische landschapsvorming in Binnen-Vlaanderen. Bulletin de la Société belge d'Études géographiques 32, 1963, 329-433.
- De Moor/Heyse 1978: G. De Moor / I. Heyse, De morfologische evolutie van de Vlaamse Vallei. De Aardrijkskunde 4, 1978, 343-375.
- Denys/Kiden/Verbruggen 1998: L. Denys / P. Kiden / C. Verbruggen, Opaline concretions in Weichselian Late-glacial lake marl from Flanders, northern Belgium. Journal of Paleolimnology 20, 1998, 91-98.
- Denys/Verbruggen/Kiden 1990: L. Denys / C. Verbruggen / P. Kiden, Palaeolimnological aspects of a Late-Glacial shallow lake in Sandy Flanders, Belgium. Hydrobiologia 214, 1990, 173-278.
- Derese et al. 2010: C. Derese / D. A. G. Vandenberghe / A. Zwertvaegher / J. Verniers / Ph. Crombé / P. van den Haute, The timing of aeolian events near archaeological settlements around Heidebos (Moervaart area, N Belgium). Netherlands Journal of Geosciences 89, 2010, 173-186.

- De Smedt et al. 2012: Ph. De Smedt / M. Van Meirvenne / N. S. Davies / M. Bats / T. Saey / J. De Reu / E. Meerschman / V. Gelorini / A. Zwervaegeher / Ph. Antrop / J. Bourgeois / Ph. De Maeyer / P. Finke / J. Verniers / Ph. Crombé, A multidisciplinary approach to reconstructing Late Glacial and Early Holocene landscapes. *Journal of Archaeological Science* 40, 2012, 1260-1267.
- Fagnart 1997: J.-P. Fagnart, La fin des temps glaciaires dans le Nord de la France. Approches archéologiques et environnementales des occupations humaines du Tardiglaciaire. *Mémoires de la Société préhistorique française* 24 (Paris 1997).
- Heyse 1979: I. Heyse, Bijdrage tot de geomorfologische kennis van het noordwesten van Oost-Vlaanderen (België). *Verhandelingen van de Koninklijke Academie voor Wetenschappen, Letteren en Schone Kunsten van België* 40 (Brussel 1979).
- 1983: I. Heyse, Preliminary results of the study of a Vistulian Late Glacial drainage pattern in the Scheldt basin (Belgium-Flemish Valley-Moervaart depression). *Quaternary Studies in Poland* 4, 1983, 135-143.
- Hoek/Bohncke 2002: W. Hoek / S. J. P. Bohncke, Climatic and environmental events over the Last Termination, as recorded in The Netherlands: a review. *Netherlands Journal of Geosciences* 81, 2002, 123-137.
- Kasse et al. 2005: C. Kasse / W. Z. Hoek / S. J. P. Bohncke / M. Konert / J. W. H. Weijers / M. L. Cassee / R. M. Van der Zee, Late Glacial fluvial response of the Niers-Rhine (western Germany) to climate and vegetation change. *Journal of Quaternary Science* 20, 2005, 377-394.
- Kiden 1989: P. Kiden, Temse en de Schelde: de geomorfologische achtergrond. In: H. Thoen (ed.), *Temse en de Schelde: Van IJstijd tot Romeinen* (Brussel 1989) 13-27.
- 1991: P. Kiden, The Late-glacial and Holocene evolution of the Middle and Lower River Scheldt, Belgium. In: L. Starkel / K. L. Gregory / J. B. Thornes (eds), *Temperate Palaeohydrology* (London 1991) 283-299.
- Louwagie/Langohr 2005: G. Louwagie / R. Langohr, Palaeo-environment. *Pedo-lithostratigraphical analyses*. In: Crombé 2005, 27-107.
- Meylemans et al. 2013: E. Meylemans / F. Bogemans / A. Storme / Y. Perdaen / I. Verdurmen / K. Deforce, Lateglacial and Holocene fluvial dynamics in the Lower Scheldt basin (N-Belgium) and their impact on the presence, detection and preservation potential of the archaeological record. *Quaternary International* 308/309, 2013, 148-161.
- Missiaen et al. 2015: T. Missaen / J. Verhegge / K. Heirman / Ph. Crombé, Potential of Cone Penetrating Testing for mapping deeply buried palaeolandscapes in the context of archaeological surveys in polder areas. *Journal of Archaeological Science* 55, 2015, 174-187.
- Parent/Van der Plaetsen/Vanmoerkerke 1986/1987: J.-P. Parent / P. Van der Plaetsen / J. Vanmoerkerke, Prehistorische jagers en veetelers aan de Donk te Oudenaarde. *VOBOV-info* 24/25 (Gent 1986/1987).
- Perdaen/Ryssaert 2002: Y. Perdaen / C. Ryssaert, The Final Palaeolithic site of Verrebroek Dok 2: a confrontation between debitage-typology, attribute analysis and refitting. *Notae Praehistoricae* 22, 2002, 75-81.
- Perdaen/Crombé/Sergant 2004: Y. Perdaen / Ph. Crombé / J. Sergant, Federmessergroepen in de Scheldepolders (Oost-Vlaanderen, België): technologische variabiliteit binnen een vast schema. *Notae Praehistoricae* 24, 2004, 81-88.
- Perdaen/Meylemans/Vanholme 2013: Y. Perdaen / E. Meylemans / N. Vanholme, Preventief onderzoek in het kader van het Sigma-plan te Wichelen-Bergenmeersen (Oost-Vlaanderen, B). *Notae Praehistoricae* 33, 2013, 75-89.
- Perdaen et al. 2011a: Y. Perdaen / E. Meylemans / F. Bogemans / K. Deforce / A. Storme / I. Verdurmen, Op zoek naar prehistorische resten in de wetlands van de Sigmacluster Kalkense Meersen. Prospectief en evaluerend archeologisch onderzoek in het gebied Wijmeers 2, zone D/E (Wichelen, prov. Oost-Vl.). *Relicta* 8, 2011, 9-46.
- 2011b: Y. Perdaen / J. Sergant / E. Meylemans / A. Storme / K. Deforce / J. Bastiaens / S. Debruyne / A. Eryvynck / R. Langohr / A. Lentacker / K. Haneca / E. Du Rang / Ph. Crombé, Noodonderzoek van een wetland site in Bazel-Sluis (Kruibeke, Oost-Vlaanderen, B): een nieuwe kijk op de neolithisatie in Vlaanderen. *Notae Praehistoricae* 31, 2011, 31-45.
- Reimer et al. 2013: P. J. Reimer / E. Bard / A. Bayliss / J. W. Beck / P. G. Blackwell / C. Bronk Ramsey / C. E. Buck / H. Cheng / R. L. Edwards / M. Friedrich / P. M. Grootes / T. P. Guilderson / H. Hafflidason / I. Hajdas / C. Hatté / T. J. Heaton / D. L. Hoffmann / A. G. Hogg / K. A. Hughen / K. F. Kaiser / B. Kromer / S. W. Manning / M. Niu / R. W. Reimer / D. A. Richards / E. M. Scott / J. R. Southon / R. A. Staff / C. S. M. Turney / J. van der Plicht, *IntCal13 and Marine13 radiocarbon age calibration curves 0-50,000 years cal BP*. *Radiocarbon* 55, 2013, 1869-1887.
- Sergant et al. 2016: J. Sergant / H. Vandendriessche / G. Nogens / F. Cruz / L. Allemeersch / K. Aluwé / J. Jacobs / F. Wuyts / S. Windey / J. Rozek / I. Depaepe / D. Herremans / P. Laloo / Ph. Crombé, Opgraving van een mesolithische wetlandsite te Kerkhove »Stuw« (Avelgem, West-Vlaanderen): eerste resultaten. *Notae Praehistoricae* 36, 2016, 47-57.
- Storme et al. 2017: A. Storme / S. Louwye / Ph. Crombé / K. Deforce, Postglacial evolution of vegetation and environment in the Scheldt Basin (northern Belgium). *Vegetation History and Archaeobotany* 26, 2017, 293-311.
- Street et al. 2006: M. Street / F. Gelhausen / S. Grimm / F. Moseler / L. Niven / M. Sensburg / E. Turner / S. Wenzel / O. Jöris, L'occupation du bassin de Neuwied (Rhénanie centrale, Allemagne) par les Magdaléniens et les groupes à Federmesser (aziliens). *Bulletin de la Société préhistorique française* 103, 2006, 753-780.
- Tavernier/De Moor 1974: R. Tavernier / G. De Moor, L'évolution du bassin de l'Escaut. In: P. Macar (ed.), *L'évolution quaternaire des bassins fluviaux de la Mer du Nord Méridionale*. *Centenaire de la Société Géologique de Belgique* 1974, 159-231.
- Vandendriessche et al. 2016: H. Vandendriessche / A. Verbrugge / A. De Graeve / B. Cherretté / Ph. Crombé, De midden- en final-paleolithische vondsten van Ronse-Pont West (Oost-Vlaanderen, BE). *Notae Praehistoricae* 36, 2016, 91-102.
- Van Huissteden/Kasse 2001: J. Van Huissteden / C. Kasse, Detection of rapid climate change in Last Glacial fluvial successions in The Netherlands. *Global and Planetary Change* 28, 2001, 319-339.
- Vanmaercke-Gottigny 1964: M. C. Vanmaercke-Gottigny, La géomorphologie de l'Escaut d'Oudenaarde. *Acta Geographica Lovaniensia* 3, 1964, 443-473.
- Vanmoerkerke 1986: J. Vanmoerkerke, A Neolithic site at Asper-Jolleveld. *Scholae Archaeologicae* 5, 1986, 161-167.

- 1988: J. Vanmoerkerke, Een Midden-Neolithische site te Spiere. Archeologische en Historische monografieën van Zuid-West-Vlaanderen 19 (Kortrijk 1988).
- Vanmoerkerke/De Belie 1984: J. Vanmoerkerke / A. De Belie, Epipaleolithicum en laat-neolithicum te Klein-Sinaai (Stekene). VO-BOV-info 14, 1984, 1-13.
- Vanmontfort et al. 2001/2002: B. Vanmontfort / A.-I. Geerts / Ch. Casseyas / C. Bakels / Ch. Buydens / F. Damblon / R. Langohr / W. Van Neer / P. M. Vermeersch, De Hel in de tweede helft van het 5de millennium v. Chr. Een midden-Neolithische enclosure te Spiere (prov. West-Vlaanderen). Archeologie in Vlaanderen 8, 2001/2002, 9-77.
- Van Strydonck 2005: M. Van Strydonck, Palaeo-environment. Radiocarbon dating. In: Crombé 2005, 127-137.
- Van Strydonck/Crombé 2005: M. Van Strydonck / Ph. Crombé, Features: radiocarbon dating. In: Crombé 2005, 180-212.
- Verbruggen 1971: C. Verbruggen, Postglaciale landschapsgeschiedenis van Zandig Vlaanderen [unpubl. Diss. Univ. Ghent 1971].
- Verbruggen/Denys/Kiden 1996: C. Verbruggen / L. Denys / P. Kiden, Belgium. In: B. E. Berglund / H. J. B. Birks / M. Ralska-Jasiewiczowa / H. E. Wright (eds), Palaeoecological Events during the Last 15 000 Years: Regional Syntheses of Palaeoecological Studies of Lakes and Mires in Europe (Chichester 1996) 553-574.
- Verhegge/Missiaen/Crombé 2016: J. Verhegge / T. Missiaen / Ph. Crombé, Exploring integrated geophysics and geotechnics as a palaeolandscape reconstruction tool: archaeological prospection of (prehistoric) sites buried deeply below the Scheldt polders (NW Belgium). Archaeological Prospection 23, 2016, 125-145.
- Vermeersch 1976: P. M. Vermeersch, Un site Tjongerien à Harelbeke Gavermeersen. Westvlaamse Archaeologica 4, 1976, 33-71.

Summary

In this paper the spatial distribution of Federmesser-Gruppen (FMG) sites in the Scheldt Basin is discussed. A clear pattern of focused occupation and exploitation of the dry banks of former dune slacks and freshwater lakes is observed, while the river floodplains were seemingly avoided. The latter may be due to the highly dynamic environment of the rivers, which were characterised by very high discharges and continuous erosion. During the late Allerød (GI-1a) and the transition towards the Younger Dryas (GS-1) the meandering rivers finally stabilised, gradually turning into slow running and shallow streams. This probably was the result of a general decrease of the ground water table, which also affected the inland lakes and ponds, reducing them to dry or swampy depressions. It is assumed that this marked hydrological event, which may be linked to cooling during the Intra Allerød Cold Period (GI-1b) and the subsequent Younger Dryas (GS-1), had a considerable impact on the settlement system of the FMG hunter-gatherers, who may have been forced to leave the Scheldt Basin.

Keywords

Scheldt Basin, Federmesser-Gruppen, site distribution, hydrological change, climate change